









a Find the modal class <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> $20 < x \leq 30$ the modal class	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	b Which class contains the median? <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> $10 < x \leq 20$ the class with the median	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	c Estimate the mean <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> 16.5 an estimated mean from midpoints	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	d Find the lower quartile of 2, 5, 6, 9, 11, 14, 15 5 the lower quartile
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
e Find the interquartile range of 13, 4, 8, 17, 6, 10, 3 9 the interquartile range: order first	f Find the interquartile range of 3, 4, 6, 7, 8, 10, 12, 13, 16, 18, 20 10 quartiles of eleven values	g Find the median  20 the median from a box plot	h Find the range  40 the range from a box plot																														
i Find the interquartile range  15 the IQR from a box plot	j What fraction of the data is above the upper quartile? $\frac{1}{4}$ a quarter above the upper quartile	k Which is more consistent, A or B?  A (smaller IQR) comparing spread with the IQR	l A data set has 80 values. How many are between the quartiles? 40 half the data between the quartiles																														

a Find the modal class <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> $20 < x \leq 30$ the modal class	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	b Which class contains the median? <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> $10 < x \leq 20$ the class with the median	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	c Estimate the mean <table border="1"> <thead> <tr> <th>Class</th> <th>Frequency</th> </tr> </thead> <tbody> <tr> <td>$0 < x \leq 10$</td> <td>7</td> </tr> <tr> <td>$10 < x \leq 20$</td> <td>4</td> </tr> <tr> <td>$20 < x \leq 30$</td> <td>8</td> </tr> <tr> <td>$30 < x \leq 40$</td> <td>1</td> </tr> </tbody> </table> 16.5 an estimated mean from midpoints	Class	Frequency	$0 < x \leq 10$	7	$10 < x \leq 20$	4	$20 < x \leq 30$	8	$30 < x \leq 40$	1	d Find the lower quartile of 2, 5, 6, 9, 11, 14, 15 5 the lower quartile
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
Class	Frequency																																
$0 < x \leq 10$	7																																
$10 < x \leq 20$	4																																
$20 < x \leq 30$	8																																
$30 < x \leq 40$	1																																
e Find the interquartile range of 13, 4, 8, 17, 6, 10, 3 9 the interquartile range: order first	f Find the interquartile range of 3, 4, 6, 7, 8, 10, 12, 13, 16, 18, 20 10 quartiles of eleven values	g Find the median  20 the median from a box plot	h Find the range  40 the range from a box plot																														
i Find the interquartile range  15 the IQR from a box plot	j What fraction of the data is above the upper quartile? $\frac{1}{4}$ a quarter above the upper quartile	k Which is more consistent, A or B?  A (smaller IQR) comparing spread with the IQR	l A data set has 80 values. How many are between the quartiles? 40 half the data between the quartiles																														